

2018

MATHEMATICS

(Major)

Paper : 6.2

(Numerical Analysis)

Full Marks : 60

Time : 3 hours

*The figures in the margin indicate full marks
for the questions*

1. Answer the following questions :

1×7=7

- (a) What do you mean by 'normalized floating point representation' of real numbers?
- (b) Define 'round-off' error.
- (c) Write down the approximate representation of $\frac{2}{3}$ correct to four significant figures and find the absolute error.

(2)

(d) Give the relationship between the operators E and Δ .

(e) Evaluate $E^2 x^2$ when $h=2$.

(f) Show that $E \nabla \equiv \nabla E \equiv \Delta$.

(g) Write the general quadrature formula in numerical integration.

2. Answer the following questions : $2 \times 4 = 8$

(a) Explain briefly the importance of numerical differentiation method.

(b) Determine the number of significant figures in 8.1205 given its absolute error as 0.3×10^{-2} .

(c) Evaluate :

$$\left(\frac{\Delta^2}{E} \right) x^3$$

(d) Write the numerical differentiation formulae for finding the first and second derivatives of a function $f(x)$ at a point x near the beginning of a given set of tabulated values.

8A/902

(Continued)

(3)

3. Answer the following questions : $5 \times 3 = 15$

(a) Find the absolute, relative and percentage error when $\frac{3}{7}$ is approximated by 0.4286.

(b) Using the method of separation of symbols, prove that

$$u_x = u_{x-1} + \Delta u_{x-2} + \Delta^2 u_{x-3} + \dots + \Delta^{n-1} u_{x-n} + \Delta^n u_{x-n}$$

$$\text{where } u_{x+h} = E^h u_x.$$

Or

In an examination, the number of candidates who obtained marks between certain limits were as follows :

Marks	No. of candidates
0-19	41
20-39	62
40-59	65
60-79	50
80-99	17

Find the number of candidates who obtained less than 70 marks using a suitable interpolation formula.

(c) Find the polynomial of the lowest possible degree which assumes the values 3, 12, 15, -21 when x takes the values 3, 2, 1, -1 respectively.

8A/902

(Turn Over)

(4)

Or

Construct Lagrange's interpolating polynomial using the following data :

x	: 40	45	50	55
$y = f(x)$: 15.22	13.99	12.62	11.13

4. Answer any one part :

- (a) (i) Given $u_{20} = 24$, $u_{24} = 32$, $u_{28} = 35$, $u_{32} = 40$, find u_{25} by Bessel's formula.

- (ii) Find $f'(4)$ and $f''(4)$ from the following data :

x	: 1	2	4	8	10
$f(x)$: 0	1	5	21	27

5+5=10

- (b) (i) Use Stirling's formula to find a polynomial of degree three or less which takes the following values of the function u_x :

x	: 4	6	8	10
u_x	: 1	3	8	20

- (ii) Find the value of

$$\int_0^1 \frac{x^2}{1+x^3} dx$$

using Simpson's $\frac{1}{3}$ rd rule, dividing the range into four equal parts. 5+5=10

8A/902

(Continued)

(5)

5. Answer any one part :

- (a) (i) Evaluate

$$\int_4^{5.2} \log_e x dx$$

by Weddle's rule. Also compute the error.

- (ii) Derive Simpson's $\frac{1}{3}$ rd rule from Newton-Cotes quadrature formula.

5+5=10

- (b) (i) A curve is drawn to pass through the points given by the following table :

x	: 1	1.5	2	2.5	3	3.5	4
y	: 2	2.4	2.7	2.8	3	2.6	2.1

Estimate the area bounded by the curve, the x -axis and the lines $x = 1$ and $x = 4$.

- (ii) The following table gives the values of acceleration f of a particle in cm/sec^2 at equal interval of time t in sec.

t	: 0.0	0.5	1.0	1.5	2.0
f	: 0.3989	0.3521	0.2420	0.1295	0.0540

Find the velocity of the particle at $t = 2$ sec.

5+5=10

8A/902

(Turn Over)

6. Answer any one part :

(a) (i) Derive the rate of convergence of the secant method.

(ii) Use Newton-Raphson method to find a positive root of the equation $e^x - 3x = 0$ correct to four decimal places. 5+5=10

(b) (i) Find an approximate real root of the equation $x^3 - x - 1 = 0$ using bisection method. Perform four iterations.

(ii) Using Regula-Falsi method, find the approximate real root of the equation

$$x \log_{10} x - 1.2 = 0$$

correct to five decimal places. 5+5=10
