

2019

MATHEMATICS

( Major )

Paper : 5.1

( Real and Complex Analysis )

Full Marks : 60

Time : 3 hours

The figures in the margin indicate full marks  
for the questions

Symbols have usual meaning

1. Answer the following questions :

1×7=7

- (a) Write down a sufficient condition for the equality of  $f_{xy}$  and  $f_{yx}$ .
- (b) Give an example of a discontinuous function which is Riemann integrable.
- (c) If  $P^*$  is a refinement of a partition  $P$  of a bounded function  $f$ , then write down the relations between  $U(P, f)$ ,  $U(P^*, f)$ ,  $L(P, f)$ ,  $L(P^*, f)$ .

( 2 )

(d) Define pole of order  $n$  of a complex valued function  $f(z)$ .

(e) A function  $f(z) = u(x, y) + iv(x, y)$  is defined such that  $\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}$ . State whether  $f$  is analytic or not.

(f) Let  $f(z) = u(x, y) + iv(x, y)$  be analytic in a region  $R$ . Prove that  $\frac{\partial(u, v)}{\partial(x, y)} = |f'(z)|^2$ .

(g) Find the fixed points of the transformation  $w = z + 5$ .

2. Answer the following questions : 2×4=8

(a) Show that

$$\lim_{x \rightarrow 0} \lim_{y \rightarrow 0} f(x, y) = \lim_{y \rightarrow 0} \lim_{x \rightarrow 0} f(x, y)$$

but  $\lim_{(x, y) \rightarrow (0, 0)} f(x, y)$  does not exist, where

$$f(x, y) = \frac{xy}{x^2 + y^2}, \quad (x, y) \neq (0, 0)$$

$$= 0, \quad (x, y) = (0, 0)$$

20A/269

( Continued )

( 3 )

(b) Prove that the improper integral

$$\int_a^b \frac{dx}{(x-a)^n}$$

converges if and only if  $n < 1$ .

(c) Let  $C$  be the curve in the  $xy$ -plane defined by  $3x^2y - 2y^3 = 5x^4y^2 - 6x^2$ . Find a unit vector normal to  $C$  at  $(1, -1)$ .

(d) Show that

$$\nabla \equiv \frac{\partial}{\partial x} + i \frac{\partial}{\partial y} = 2 \frac{\partial}{\partial \bar{z}}$$

3. Answer any three parts :

$$5 \times 3 = 15$$

(a) Show that the function

$$f(x, y) = \frac{x^2y}{x^4 + y^2}, \quad x^2 + y^2 \neq 0$$

$$= 0, \quad x = y = 0$$

possesses first partial derivatives everywhere, including the origin, but the function is discontinuous at the origin.

(b) Prove that a bounded function  $f$  is integrable on  $[a, b]$  iff for every  $\epsilon > 0$ , there exists a partition  $P$  of  $[a, b]$  such that  $U(P, f) - L(P, f) < \epsilon$ .

20A/269

( Turn Over )

( 4 )

(c) Prove that every absolutely convergent improper integral is convergent.

(d) Given,  $u = e^{-x}(x \sin y - y \cos y)$ , find  $v$  such that  $f(z) = u + iv$  is analytic.

(e) Evaluate  $\int_C \bar{z} dz$  from  $z = 0$  to  $z = 4 + 2i$  along the curve  $C$  given by (i)  $z = t^2 + it$  and (ii) the line from  $z = 0$  to  $z = 2i$  and then the line from  $z = 2i$  to  $z = 4 + 2i$ .

4. Answer any one part :

10

(a) (i) Show that  $f(xy, z - 2x) = 0$ ,  $f$  is differentiable and  $f_v \neq 0$ , where  $v = z - 2x$  satisfies the equation

$$x \frac{\partial z}{\partial x} - y \frac{\partial z}{\partial y} = 2x$$

5

(ii) Show that the function

$$f(x, y) = y^2 + x^2 y + x^4$$

has a minimum at  $(0, 0)$ .

5

( 5 )

(b) (i) Show that  $\int_0^1 x^{m-1}(1-x)^{n-1} dx$  exists if and only if  $m, n$  both are positive. 5

(ii) Show that the integral

$$\int_0^1 \frac{\sin(1/x)}{x^p} dx, \quad p > 0$$

is absolutely convergent for  $p < 1$ . 5

5. Answer any one part :

10

(a) (i) The roots of the equation in  $\lambda$

$$(\lambda - x)^3 + (\lambda - y)^3 + (\lambda - z)^3 = 0$$

are  $u, v, w$ . Prove that

$$\frac{\partial(u, v, w)}{\partial(x, y, z)} = -2 \frac{(y-z)(z-x)(x-y)}{(u-v)(v-w)(w-u)}$$

5

(ii) Prove that if  $f$  and  $g$  are Riemann integrable on  $[a, b]$ , then  $f + g, f - g$  are also Riemann integrable on  $[a, b]$ . 5

(b) (i) Show that the function  $[x]$ , where  $[x]$  denotes the greatest integer not greater than  $x$ , is Riemann integrable in  $[0, 3]$ . 5

- (ii) Prove that if a function  $f$  is bounded and integrable on  $[a, b]$  and there exists a function  $F$  such that  $F' = f$  on  $[a, b]$ , then  $\int_a^b f dx = F(b) - F(a)$ .

5

6. Answer any one part :

10

- (a) (i) Prove that if

$$w = f(z) = u(x, y) + iv(x, y)$$

is analytic, then

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y} \text{ and } \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

5

- (ii) Let  $u(x, y) = \alpha$  and  $v(x, y) = \beta$ , where  $u$  and  $v$  are the real and imaginary parts of an analytic function  $f(z)$  and  $\alpha, \beta$  are the constants, represent two families of curves. Prove that if  $f'(z) \neq 0$ , then the families are orthogonal.

5

- (b) (i) Let  $f(z)$  be analytic inside and on a circle  $C$  of radius  $r$  and centre at  $z = a$ . Then prove that

$$f^{(n)}(a) \leq \frac{Mn!}{r^n}, \quad n = 0, 1, 2, \dots$$

where  $M$  is a constant such that  $|f(z)| < M$  on  $C$  and  $f^{(n)}(a)$  represents  $n$ -th derivative of  $f(z)$  at  $z = a$ .

5

( Continued )

- (ii) Let the rectangular region  $R$  in the  $z$ -plane be bounded by  $x = 0, y = 0, x = 2, y = 1$ . Determine the region  $R'$  of the  $w$ -plane into which  $R$  is mapped under the transformation

$$1. w = z + (1 - 2i)$$

$$2. w = \sqrt{2}e^{i\pi/4}z$$

$$2+3=5$$

★ ★ ★