

Total number of printed pages-8

3 (Sem-3/CBCS) STA HC 1

2021

(Held in 2022)

**STATISTICS**

(Honours)

Paper : STA-HC-3016

**(Sampling Distributions)**

Full Marks : 60

Time : Three hours

***The figures in the margin indicate full marks for the questions.***

1. Answer the following questions as directed :

$$1 \times 7 = 7$$

- (a) For random sample of size 2 drawn from  $N(0, \sigma^2)$  population, the expected value of the smallest order statistic is

(i)  $-\frac{\sigma^2}{\sqrt{\pi}}$

(ii)  $-\frac{\sigma}{\sqrt{\pi}}$

Contd.

(iii)  $\frac{2\sigma^2}{\sqrt{\pi}}$

(iv) None of the above

(Choose the correct option)

(b) If  $X$  and  $Y$  are two independent Chi-square variates with  $n_1$  and  $n_2$  degrees of freedom respectively, then  $u = \frac{X}{Y}$  follows

(i)  $\beta_2\left(\frac{n_1}{2}, \frac{n_2}{2}\right)$

(ii)  $\beta_1(n_1, n_2)$

(iii) F-distribution

(iv) None of the above

(Choose the correct option)

(c) If  $X$  is distributed as a Chi-square variate with  $n$  d.f., then for large  $N$ ,  $\sqrt{2x}$  is distributed as

(i)  $N(2n, 1)$

(ii)  $N(\sqrt{2n}, 1)$

(iii)  $N(\sqrt{2n}, n)$

(iv) None of the above

(Choose the correct option)

(d) For testing the hypothesis 'population correlation ratio is zero'. The test statistics is

(i)  $\frac{\eta^2}{1-\eta^2} \cdot \frac{N-h}{h-1}$

(ii)  $\frac{1-\eta^2}{\eta^2} \cdot \frac{N-h-1}{h}$

(iii)  $\frac{\eta^2}{1-\eta^2} \cdot \frac{N-h-1}{h^2}$

(iv) None of the above

(Choose the correct option)

(e) If a statistic  $t$  follows students  $t$ -distribution with  $n$  d.f., then  $t^2$  follows

\_\_\_\_\_. (Fill in the blank)



- (f) 95% confidence limits for population proportion are

(i)  $p \pm 1.96 \sqrt{\frac{pq}{n}}$

(ii)  $p \pm 2.58 \sqrt{\frac{pq}{n}}$

(iii)  $p \pm 2.33 \sqrt{\frac{pq}{n}}$

- (iv) None of the above

(Choose the correct option)

- (g) The moment generating function of t-distribution does not exist.

(State True or False)

2. Answer the following questions :  $2 \times 4 = 8$

- (a) Explain the terms 'level of significance' and 'critical region'.

- (b) Obtain cumulant generating function of Chi-square distribution. Hence obtain mean and variance.

- (c) A random sample of size 4 is drawn from the discrete uniform distribution

$$P(X=x) = \frac{1}{6}; x=1, 2, 3, 4, 5, 6$$

Obtain the distribution of the smallest and largest order statistic.

- (d) Let  $X_1, X_2, \dots, X_n$  be a random sample from  $N(0, 1)$ . Let us further

define  $\bar{x}_k = \frac{1}{k} \sum_{i=1}^k X_i$  and

$$\bar{X}_{n-k} = \frac{1}{n-k} \sum_{i=k+1}^n X_i$$

Find the distribution of

$$k \bar{X}_k^{-2} + (n-k) \bar{X}_{n-k}^{-2}$$

3. Answer **any three** of the following questions :  $5 \times 3 = 15$

- (a) Derive the joint probability distribution of  $X_{(r)}$  and  $W_{rs} = X_{(s)} - X_{(r)}$  ;  $(r < s)$  based on a random sample of size  $n$  from the exponential distribution with parameter  $\alpha$ .



(b) If  $X$  has a F-distribution with  $\nu_1$  and  $\nu_2$  degrees of freedom, find the distribution of  $\frac{1}{X}$ .

(c) Show that the m.g.f. of  $Y = \log \chi^2$ , where  $\chi^2$  follows Chi-square distribution with  $n$  d.f. is

$$M_Y(t) = \frac{2^t \Gamma\left(\frac{n}{2} + t\right)}{\Gamma(n/2)}$$

(d) Suppose a person is interested in testing the equality of two population standard deviations, say  $\sigma_1$  and  $\sigma_2$ . For this purpose two samples of sizes  $n_1$  and  $n_2$  are drawn from the two populations respectively and suppose that the sample standard deviations are  $S_1$  and  $S_2$  respectively.

Explain how you would test the hypothesis  $H_0: \sigma_1 = \sigma_2$ . Also discuss test of  $H_0$  when both  $n_1$  and  $n_2$  are large.

(e) Show that for large degrees of freedom,  $t$ -distribution tends to standard normal distribution.

Answer the following questions :

10×3=30

4. (a) (i) Explain clearly the procedure generally followed in testing of a hypothesis. Also point the difference between one-tail and two-tail tests. 5

(ii) Show that in odd sample of size  $n$  from  $U(0, 1)$  population, mean of the distribution of median is  $\frac{1}{2}$ . 5

**Or**

(b) Derive the probability density function of the student's  $t$ -distribution with  $\nu$  d.f. and hence find its mean and variance. 10

5. (a) (i) Show that for large d.f., the Chi-square distribution tends to the normal distribution. 5



- (ii) Show that for  $t$ -distribution with  $n$  d.f. mean deviation about mean is given by

$$\frac{\sqrt{n} \Gamma\left(\frac{n-1}{2}\right)}{\sqrt{\pi} \Gamma\left(\frac{n}{2}\right)}$$

5

6. (a) (i) Derive the expression for the standard error of the mean of a random sample of size  $n$  and sample proportion. 5

- (ii) Write down some familiar applications of order statistics. 5

**Or**

- (b) (i) If  $n_2 \rightarrow \infty$  in  $F(n, n_2)$  distribution, then show that  $\chi^2 = n_1 F$  follows Chi-square distribution with  $n_1$  d.f. 7

- (ii) Explain how the student's  $t$ -distribution is used to test the difference between the means of two samples which are paired together. 3