2022

MATHEMATICS

(Major)

Paper: 5.5

(Probability)

Full Marks: 60

Time: 3 hours

The figures in the margin indicate full marks for the questions

- 1. Answer the following questions as directed: 1×8=8
 - (a) If A and B are mutually exclusive, what will be the modified statement of $P(A \cup B) = P(A) + P(B) P(A \cap B)$?

- (b) If a non-negative real valued function f is the probability density function of some continuous random variable, then what is the value of $\int_{-\infty}^{\infty} f(x) dx$?
- (c) What is meant by mathematical expectation of a random variable?
- (d) For a Bernoulli random variable X with P(X = 0) = 1 p and P(X = 1) = p, write E(X) and V(X) in terms of p.
- (e) Can the probabilities of three mutually exclusive events A, B, C as given by $P(A) = \frac{2}{3}$, $P(B) = \frac{1}{4}$ and $P(C) = \frac{1}{6}$ be correct? If not, give reasons.
- (f) Test the validity of the following probability distribution:

	x	-1	0	1		
	p(x)	0.4	0	1		
1107			0.4	0.3		

(g) Under what condition cov(X, Y) = 0?

- (h) Choose the correct option for binomial distribution.
 - (i) Variance = Mean
 - (ii) Variance > Mean
 - (iii) Variance < Mean
- 2. Answer the following questions: 3×4=12
 - (a) A speaks the truth in 75% cases and B in 80%. In what percentages of cases are they likely to contradict each other while narrating the same incident?
 - (b) Define probability mass function and probability density function for a random variable X.
 - (c) The second moment about any point (a) is minimum when taken about the mean (μ), i.e.,

$$E(X-a)^2 \ge E(X-\mu)^2$$

Prove or disprove the above statement.

(d) State the three Poisson's postulates.

- 3. Answer any two parts from the following questions: 5×2=10
 - (a) What is meant by partition of a sample space S? If $H_i = (i = 1, 2, ..., n)$ is a partition of the sample space S, then for any event A, prove that

$$P(H_i / A) = \frac{P(H_i) P(A/H_i)}{\sum_{i=1}^{n} P(H_i) P(A/H_i)}$$

- (b) Three machines A, B and C produce respectively 60%, 30% and 10% of the total number of items of a factory. The percentage of defective output of these machines are respectively 2%, 3% and 4%. An item is selected at random and is found to be defective. Find the probability that the item was produced by machine C.
- (c) A bag contains 5 balls and it is known how many of these are white. Two balls are drawn and are found to be white. What is the probability that all are white?

- 4. Answer any two parts from the following questions: 5×2=10
 - (a) Let F be the distribution function of a two-dimensional random variable (X, Y). If a < b and c < d, then show that

$$P(a < X \le b, c < Y \le d) = F(b,d) + F(a, c) - F(a, d) - F(b, c)$$

(b) If X is a discrete random variable having probability mass function

Mass point	0	1	2	3	4	5	6	7
P(X=x)	0	k	2k	3k	4k	k^2	$2k^2$	$7k^2 + k$

Determine (i) k, (ii)
$$P(X < 6)$$
 and (iii) $P(X \ge 6)$. $2+2+1=5$

(c) The probability density function of a two-dimensional random variable (X, Y) is given by

$$f(x, y) = x + y, \quad 0 < x + y < 1$$
$$= 0, \quad \text{elsewhere}$$

Evaluate $P(X < \frac{1}{2}, Y > \frac{1}{4})$.

- 5. Answer any two parts from the following questions: 5×2=10
- (a) From a bag containing 4 white and 6 red balls, three balls are drawn at random. Find the expected number of white balls drawn.
 - (b) For any two independent random variables X and Y, for which E(X) and E(Y) exist, show that

$$E(XY) = E(X)E(Y)$$

(c) The probability density function of a continuous bivariate distribution is given by the joint density function

$$f(x, y) = x + y$$
, $0 < x < 1$, $0 < y < 1$
= 0 , elsewhere

Find E(X), E(Y), var (X), var (Y) and E(XY).

- 6. Answer any two parts from the following questions: 5×2=10
 - (a) If X is a Poisson distributed random variable with parameter μ , then show that $E(X) = \mu$ and $var(X) = \mu$.

- (b) Show that normal distribution may be regarded as a limiting case of Poisson's distribution as the parameter $m \to \infty$.
- (c) Define binomial distribution. What is the probability of guessing correctly at least six of the ten answers in a True-False objective test?

**