

Total number of printed pages-23

3 (Sem-5/CBCS) MAT HC 1 (N/O)

2022

MATHEMATICS

(Honours)

Paper : MAT-HC-5016

(For New Syllabus)

(Complex Analysis)

Full Marks : 60

Time : Three hours

***The figures in the margin indicate
full marks for the questions.***

1. Answer **any seven** questions from the following : 1×7=7

(a) Describe the domain of definition of the

function $f(z) = \frac{z}{z + \bar{z}}$.

(b) What is the multiplicative inverse of a non-zero complex number $z = (x, y)$?

Contd.

(c) Verify that $(3, 1) (3, -1) \left(\frac{1}{5}, \frac{1}{10}\right) = (2, 1)$.

(d) Determine the accumulation points of the set $Z_n = \frac{i}{n} (n = 1, 2, 3, \dots)$.

(e) Write the Cauchy-Riemann equations for a function $f(z) = u + iv$.

(f) When a function f is said to be analytic at a point?

(g) Determine the singular points of the function $f(z) = \frac{2z+1}{z(z^2+1)}$.

(h) $\exp(2 \pm 3\pi i)$ is

(i) $-e^2$

(ii) e^2

(iii) $2e$

(iv) $-2e$ (Choose the correct answer)

(i) The value of $\log(-1)$ is

(i) 0

(ii) $2n\pi i$

(iii) πi

(iv) $-\pi i$ (Choose the correct answer)

(j) If $z = x + iy$, then $\sin z$ is

(i) $\sin x \cosh y + i \cos x \sinh y$

(ii) $\cos x \cosh y - i \sin x \sinh y$

(iii) $\cos x \sinh y + i \sin x \cosh y$

(iv) $\sin x \sinh y - i \cos x \cosh y$
(Choose the correct answer)

(k) If $\cos z = 0$, then

(i) $z = n\pi, (n = 0, \pm 1, \pm 2, \dots)$

(ii) $z = \frac{\pi}{2} + n\pi, (n = 0, \pm 1, \pm 2, \dots)$

(iii) $z = 2n\pi, (n = 0, \pm 1, \pm 2, \dots)$

(iv) $z = \frac{\pi}{2} + 2n\pi, (n = 0, \pm 1, \pm 2, \dots)$
(Choose the correct answer)

(l) If z_0 is a point in the z -plane, then

$$\lim_{z \rightarrow \infty} f(z) = \infty \text{ if}$$

$$(i) \lim_{z \rightarrow 0} \frac{1}{f(z)} = \infty$$

$$(ii) \lim_{z \rightarrow 0} f\left(\frac{1}{z}\right) = 0$$

$$(iii) \lim_{z \rightarrow 0} \frac{1}{f(z)} = 0$$

$$(iv) \lim_{z \rightarrow 0} \frac{1}{f\left(\frac{1}{z}\right)} = 0$$

(Choose the correct answer)

2. Answer **any four** questions from the following :

$$2 \times 4 = 8$$

(a) Reduce the quantity $\frac{5i}{(1-i)(2-i)(3-i)}$ to a real number.

(b) Define a connected set and give *one* example.

(c) Find all values of z such that $\exp(2z-1) = 1$.

(d) Show that $\log(i^3) \neq 3 \log i$.

(e) Show that

$$2 \sin(z_1 + z_2) \sin(z_1 - z_2) = \cos 2z_2 - \cos 2z_1$$

(f) If z_0 and w_0 are points in the z plane and w plane respectively, then prove that $\lim_{z \rightarrow z_0} f(z) = \infty$ if and only if

$$\lim_{z \rightarrow z_0} \frac{1}{f(z)} = 0.$$

(g) State the Cauchy integral formula. Find

$\frac{1}{2\pi i} \int_C \frac{1}{z-z_0} dz$ if z_0 is any point interior to simple closed contour C .

(h) Show that $\int_0^{\frac{\pi}{6}} e^{i2t} dt = \frac{\sqrt{3}}{4} + \frac{i}{4}$.

3. Answer **any three** questions from the following : 5×3=15

(a) (i) If a and b are complex constants, use definition of limit to show that

$$\lim_{z \rightarrow z_0} (az + b) = az_0 + b. \quad 2$$

(ii) Show that

$$\lim_{z \rightarrow 0} \left(\frac{z}{\bar{z}} \right)^2 \text{ does not exist.} \quad 3$$

(b) Suppose that $\lim_{z \rightarrow z_0} f(z) = w_0$ and

$$\lim_{z \rightarrow z_0} F(z) = W_0.$$

Prove that $\lim_{z \rightarrow z_0} [f(z)F(z)] = w_0W_0$.

(c) (i) Show that for the function $f(z) = \bar{z}$, $f'(z)$ does not exist anywhere. 3

(ii) Show that $\lim_{z \rightarrow \infty} \frac{4z^2}{(z-1)^2} = 4. \quad 2$

(d) (i) Show that the function $f(z) = \exp \bar{z}$ is not analytic anywhere. 3

(ii) Find all roots of the equation

$$\log z = i \frac{\pi}{2}. \quad 2$$

(e) If a function f is analytic at all points interior to and on a simple closed contour C , then prove that

$$\int_C f(z) dz = 0.$$

(f) Evaluate : 2½+2½=5

$$(i) \int_C \frac{e^{-z}}{z - (\pi i/2)} dz$$

$$(ii) \int_C \frac{z}{2z+1} dz$$

where C denotes the positively oriented boundary of the square whose sides lie along the lines $x = \pm 2$ and $y = \pm 2$.

(g) Prove that any polynomial

$$P(z) = a_0 + a_1z + a_2z^2 + \dots + a_nz^n \quad (a_n \neq 0)$$

of degree n ($n \geq 1$) has at least one zero.

(h) Find the Laurent series that represents

$$\text{the function } f(z) = z^2 \sin\left(\frac{1}{z^2}\right) \text{ in the}$$

domain $0 < |z| < \infty$.

4. Answer **any three** questions from the following : $10 \times 3 = 30$

(a) (i) If a function f is continuous throughout a region R that is both closed and bounded, then prove that there exists a non-negative real number μ such that $|f(z)| \leq \mu$ for all points z in R , where equality holds for at least one such z .

4

(ii) Let a function

$f(z) = u(x, y) + iv(x, y)$ be analytic throughout a given domain D . If $|f(z)|$ is constant throughout D , then prove that $f(z)$ must be constant there too. 3

(iii) Show that the function

$f(z) = \sin x \cosh y + i \cos x \sinh y$ is entire. 3

(b) (i) Suppose that $f(z_0) = g(z_0) = 0$ and that $f'(z_0)$ $g'(z_0)$ exist, where $g'(z_0) \neq 0$. Use definition of derivative to show that

$$\lim_{z \rightarrow z_0} \frac{f(z)}{g(z)} = \frac{f'(z_0)}{g'(z_0)}. \quad 3$$

(ii) Show that $f'(z)$ does not exist at any point if $f(z) = 2x + ixy^2$.

3

(iii) If a function f is analytic at a given point, then prove that its derivatives of all orders are analytic there too. 4

(c) Let the function

$f(z) = u(x, y) + iv(x, y)$ be defined throughout some ε -neighbourhood of a point $z_0 = x_0 + iy_0$. If u_x, u_y, v_x, v_y exist everywhere in the neighbourhood, and these partial derivatives are continuous at (x_0, y_0) and satisfy the Cauchy-Riemann equations at (x_0, y_0) , then prove that $f'(z_0)$ exist and $f'(z_0) = u_x + iv_x$ where the right hand side is to be evaluated at (x_0, y_0) .

Use it to show that for the function $f(z) = e^{-x} \cdot e^{-y}$, $f''(z)$ exists everywhere and $f''(z) = f(z)$. 6+4=10

(d) (i) Prove that the existence of the derivative of a function at a point implies the continuity of the function at that point.

With the help of an example show that the continuity of a function at a point does not imply the existence of derivative there.

3+5=8

(ii) Find $f'(z)$ if

$$f(z) = \frac{z-1}{2z+1} \left(z \neq -\frac{1}{2} \right). \quad 2$$

(e) (i) Prove that $\int_C \frac{dz}{z} = \pi i$ where C is

the right-hand half $z = 2e^{i\theta}$

$\left(-\frac{\pi}{2} \leq \theta \leq \frac{\pi}{2} \right)$ of the circle $|z| = 2$

from $z = -2i$ to $z = 2i$. 5

(ii) If a function f is analytic everywhere inside and on a simple closed contour C , taken in the positive sense, then prove that

$$f'(z) = \frac{1}{2\pi i} \int_C \frac{f(s)}{(s-z)^2} ds \quad \text{where } s$$

denotes points on C and z is interior to C . 5

(f) (i) Evaluate $I = \int_C z^{a-1} dz$

where C is the positively oriented circle $z = Re^{i\theta}$ ($-\pi \leq \theta \leq \pi$) about the origin and a denote any non-zero real number.

If a is a non-zero integer n , then what is the value of $\int_C z^{n-1} dz$?

$$4+1=5$$

(ii) Let C denote a contour of length L , and suppose that a function $f(z)$ is piecewise continuous on C . If μ is a non-negative constant such that $|f(z)| \leq \mu$ for all point z on C at which $f(z)$ is defined, then prove

$$\text{that } \left| \int_C f(z) dz \right| \leq \mu L.$$

$$\text{Use it to show that } \left| \int_C \frac{dz}{z^2 - 1} \right| \leq \frac{\pi}{3}$$

where C is the arc of the circle $|z| = 2$ from $z = 2$ to $z = 2i$ that lies in the 1st quadrant.

$$3+2=5$$

(g) (i) Apply the Cauchy-Goursat theorem to show that $\int_C f(z) = 0$ when the contour C is the unit circle $|z| = 1$, in either direction and $f(z) = ze^{-z}$. 4

(ii) If C is the positively oriented unit circle $|z| = 1$ and $f(z) = \exp(2z)$ find $\int_C \frac{f(z)}{z^4} dz$. 3

(iii) Let z_0 be any point interior to a positively oriented simple closed curve C . Show that

$$\int_C \frac{dz}{(z - z_0)^{n+1}} = 0, (n = 1, 2, \dots). \quad 3$$

(h) (i) Suppose that $z_n = x_n + iy_n$, ($n = 1, 2, \dots$) and $z = x + iy$. Prove that $\lim_{n \rightarrow \infty} z_n = z$ if and only if

$$\lim_{n \rightarrow \infty} x_n = x \text{ and } \lim_{n \rightarrow \infty} y_n = y. \quad 5$$

(ii) Show that

$$z^2 e^{3z} = \sum_{n=2}^{\infty} \frac{3^{n-2}}{(n-2)!} z^n \quad (|z| < \infty) \quad 5$$